**QUEEN'S
UNIVERSITY
BELFAST**

Influence of the cation on the solubility of CO₂ and H₂ in ionic liquids based on the bis(trifluoromethylsulfonyl)imide anion

Jacquemin, J., Husson, P., Majer, V., & Costa Gomes, M. F. (2007). Influence of the cation on the solubility of CO₂ and H₂ in ionic liquids based on the bis(trifluoromethylsulfonyl)imide anion. *Journal of Solution Chemistry*, 36(8), 967-979. <http://www.scopus.com/inward/record.url?eid=2-s2.0-34547265659&partnerID=40&md5=b8af5cc9a50e03c53aeb05523c7c2d12>

Published in:

Journal of Solution Chemistry

Document Version:

Early version, also known as pre-print

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Influence of the Cation on the Solubility of CO₂ and H₂ in Ionic Liquids Based on the Bis(trifluoromethylsulfonyl)imide Anion

Johan Jacquemin · Pascale Husson · Vladimir Majer · Margarida F. Costa Gomes

Received: 13 October 2006 / Accepted: 20 February 2007 /

Published online: 8 June 2007

© Springer Science+Business Media, LLC 2007

Abstract Experimental values for the solubility of carbon dioxide and hydrogen in three room temperature ionic liquids based on the same anion—(bistrifluoromethylsulfonyl)imide [Ntf₂][−]—and three different cations—1-butyl-3-methylimidazolium, [C₄mim], 1-ethyl-3-methylimidazolium, [C₂mim] and trimethyl-butylammonium, [N₄₁₁₁]⁺—are reported between 283 and 343 K and close to atmospheric pressure. Carbon dioxide, with a mole-fraction solubility of the order of 10^{−2}, is two orders of magnitude more soluble than hydrogen. The solubility of CO₂ is very similar in the three ionic liquids although slightly lower in the presence of the [C₂mim] cation. In the case of H₂, noticeable differences were observed with larger mole fraction solubilities in the presence of [N₄₁₁₁]⁺ followed by [C₄mim]. All of the mole-fraction solubilities decrease with increasing temperature. From the variation of Henry's law constants with temperature, the thermodynamic functions of solvation were calculated. The precision of the experimental data, considered as the average absolute deviation of the Henry's law constants from appropriate smoothing equations, is always better than ±1%.

Keywords Solubility · Carbon dioxide · Hydrogen · Ionic liquids · Ntf₂

1 Introduction

In the present paper we report original low-pressure solubility measurements of two gases in three ionic liquids. The main objective of this project is to identify the effect of cation substitution of the ionic liquid on gas solubility.

Knowledge of the gas solubility in this new family of liquid solvents is useful both from practical and fundamental points of view. Gas solubility data are valuable in view of the use of room temperature ionic liquids as reaction systems or in separation processes. They also

J. Jacquemin · P. Husson · V. Majer · M.F. Costa Gomes (✉)
Laboratoire de Thermodynamique des Solutions et des Polymères, UMR 6003 CNRS/Université
Blaise Pascal, Clermont-Ferrand, 24 avenue des Landais, 63177, Aubière Cedex, France
e-mail: margarida.c.gomes@univ-bpclermont.fr

provide a reliable way to calculate the thermodynamic properties of a solution and to assess the molecular mechanisms involved in the solvation process.

Three ionic liquids were chosen for this study: 1-ethyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide, $[\text{C}_2\text{mim}][\text{Ntf}_2]$; 1-butyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide, $[\text{C}_4\text{mim}][\text{Ntf}_2]$; and butyltrimethylammonium bis(trifluoromethylsulfonyl)imide, $[\text{N}_{4111}][\text{Ntf}_2]$.

The solubility of gases in 1-alkyl-3-methyl-imidazolium bis(trifluoromethylsulfonyl)imide has been determined by other research groups using different techniques, and Henry's law constants could be determined both from data obtained at pressures close to atmospheric [1–4] or by extrapolation of values obtained at higher pressures [5–9]. Carbon dioxide is the most studied gaseous solute and the values reported in the literature for the Henry's law constants agree to within $\pm 20\%$. Hydrogen solubility was measured only by Dyson et al. [5] at 298 K. No results are available in literature for gas solubilities in ionic liquids based on the ammonium cation studied in this work.

In this paper, experimental solubilities of carbon dioxide and hydrogen in three ionic liquids sharing the $[\text{Ntf}_2]$ anion are reported as a function of temperature between 283 and 343 K near 0.1 MPa using a high precision isochoric saturation method [10–13]. These solubility data allow the calculation of the Henry's law constants as a function of temperature, and from them the thermodynamic properties of solvation. The Gibbs energy of solvation, considered as the change in Gibbs energy when the solute is transferred at constant temperature from a pure perfect gas at 0.1 MPa to the infinitely dilute standard state in the solvent, is calculated directly from the Henry's law constants, and the standard enthalpy and entropy of solvation from their variation with temperature.

2 Experimental

2.1 Materials

The samples of the three selected ionic liquids based on the $[\text{Ntf}_2]$ anion (minimum stated purity of 0.99 in mole fraction) were supplied by the group of Prof. P. Wasserscheid (University of Erlangen-Nürnberg, Germany). The $[\text{C}_2\text{mim}][\text{Ntf}_2]$ and $[\text{C}_4\text{mim}][\text{Ntf}_2]$ were synthesized via ion exchange from $[\text{C}_2\text{mim}][\text{Br}]$ and $[\text{C}_4\text{mim}][\text{Cl}]$, respectively. Halide content analysis indicated values lower than 50 ppm for both bromide and chloride. The ionic liquid $[\text{N}_{4111}][\text{Ntf}_2]$ did not contain any chloride or bromide ions.

All samples were kept under vacuum for 15 h at 303 K before each measurement. A volumetric Karl Fisher titration (Mettler Toledo DL31) was performed before and after each series of measurements and it was found that there was no variation of the quantity of water in the samples. In all cases, the water contents were approximately 50 ± 5 ppm.

The carbon dioxide and hydrogen used for this study were supplied by AGA/Linde Gaz with mole fraction purities of 0.99995 and 0.9998, respectively. All gases were used as received from the manufacturer.

2.2 Procedure

The experimental method used for these gas solubility measurements is based on an isochoric saturation technique that has been described in previous publications [10–13]. In this technique, a known quantity of a gaseous solute is put into contact with a precisely determined quantity of degassed solvent at a constant temperature inside a cell of accurately

known volume. When thermodynamic equilibrium is attained, the pressure above the liquid solution is constant and is directly related to the solubility of the gas in the liquid.

The quantity of ionic liquid introduced into the equilibration cell, V_{liq} , is determined gravimetrically. The amount of solute present in the liquid solution, n_2^{liq} (subscript 2 stands for solute and subscript 1 stands for solvent), is calculated by the difference between two pVT measurements, the first when the gas is introduced into a calibrated bulb with volume V_{GB} and the second after thermodynamic equilibrium is reached:

$$n_2^{\text{liq}} = \frac{p_{\text{ini}} V_{\text{GB}}}{[Z_2(p_{\text{ini}}, T_{\text{ini}}) R T_{\text{ini}}]} - \frac{p_{\text{eq}} (V_{\text{tot}} - V_{\text{liq}})}{[Z_2(p_{\text{eq}}, T_{\text{eq}}) R T_{\text{eq}}]} \quad (1)$$

where p_{ini} and T_{ini} are the pressure and temperature in the first pVT determination, and p_{eq} and T_{eq} are the pressure and temperature at equilibrium. V_{tot} is the total volume of the equilibration cell and Z_2 is the compressibility factor for the pure gas. The solubility can then be expressed in mole fraction:

$$x_2 = \frac{n_2^{\text{liq}}}{n_1^{\text{liq}} + n_2^{\text{liq}}} \quad (2)$$

or as the Henry's law constant:

$$K_{\text{H}} = \lim_{x_2 \rightarrow 0} \frac{f_2(p, T, x_2)}{x_2} \cong \frac{\phi_2(p_{\text{eq}}, T_{\text{eq}}) p_{\text{eq}}}{x_2} \quad (3)$$

where f_2 is the fugacity of the solute and ϕ_2 its fugacity coefficient.

3 Results and Discussion

For each system studied, multiple experimental data points were obtained in the temperature interval between 283 and 343 K in steps of approximately 10 K. The experimental solubilities, expressed as mole fraction of carbon dioxide and hydrogen in the three selected ionic liquids are reported in Table 1 and Table 2. Henry's law constants can be calculated from the experimental values and are used to determine the mole fraction solubility, assuming a partial pressure of the gaseous solute equal to 0.1 MPa.

The relative atomic masses used are the ones recommended by IUPAC [14] and the values of the second virial coefficients for all the gases, necessary for the calculation of the compressibility factor, were taken from the compilation by Dymond and Smith [15]. The densities of the ionic liquid samples used as solvents during the present measurements were determined as a function of temperature with a precision of 0.01% using an Anton Paar vibrating-tube densitometer (model DMA 512 P) [16].

$$\rho_{[\text{C}_2\text{mim}][\text{Ntf}_2]}/\text{kg}\cdot\text{m}^{-3} = 1816.5 - 1001.3 \times 10^{-3}(T/\text{K}) \quad (4)$$

$$\rho_{[\text{C}_4\text{mim}][\text{Ntf}_2]}/\text{kg}\cdot\text{m}^{-3} = 1722.7 - 9549.6 \times 10^{-4}(T/\text{K}) \quad (5)$$

$$\rho_{[\text{N}_{4111}][\text{Ntf}_2]}/\text{kg}\cdot\text{m}^{-3} = 1654.9 - 8820.0 \times 10^{-4}(T/\text{K}). \quad (6)$$

Table 1 Experimental results for the solubility of H₂ in the ionic liquids expressed both as Henry's law constants, K_H , and as mole fraction, x_2 , corrected for a partial pressure of solute of 0.1 MPa, where p is the experimental equilibrium pressure and the per cent deviation is relative to the correlations of the data reported in Table 3

T/K	$p/10^2$ Pa	$K_H/10^5$ Pa	$x_2/10^{-4}$	dev %
[N ₄₁₁₁][Ntf ₂]				
282.93	790.55	1117	8.956	−0.0
292.61	816.51	1130	8.854	−0.0
303.35	461.05	1169	8.562	+0.4
303.38	845.38	1175	8.518	−0.1
303.40	826.75	1175	8.516	−0.2
313.29	871.97	1239	8.078	−0.1
313.29	475.04	1232	8.122	+0.5
323.16	898.44	1327	7.543	−0.1
323.18	489.00	1330	7.521	−0.4
323.24	878.71	1328	7.534	−0.2
333.07	925.01	1438	6.960	−0.2
333.27	904.94	1431	6.991	+0.4
343.02	925.01	1570	6.374	+0.0
[C ₄ mim][Ntf ₂]				
283.45	761.03	1274	7.855	−0.0
293.30	748.70	1328	7.537	−0.2
293.41	786.80	1334	7.502	+0.2
303.24	812.22	1427	7.013	+0.3
303.42	773.63	1422	7.038	−0.2
313.31	838.26	1551	6.450	−0.2
323.25	822.43	1735	5.767	+0.7
323.28	864.01	1717	5.829	−0.4
333.12	889.40	1925	5.197	−0.3
343.08	915.05	2181	4.587	−0.4
343.13	871.18	2201	4.545	+0.5
[C ₂ mim][Ntf ₂]				
282.98	806.80	1351	7.404	+0.2
293.10	834.59	1428	7.006	−0.4
303.38	801.90	1534	6.524	−0.0
303.43	433.75	1535	6.519	−0.1
304.35	865.47	1539	6.500	+0.4
313.19	826.88	1677	5.966	+0.1
313.28	446.86	1685	5.939	−0.3
323.18	460.01	1867	5.358	+0.1
325.15	857.35	1910	5.239	+0.1
333.03	877.36	2104	4.756	−0.0
333.09	473.13	2103	4.758	+0.1
343.05	902.80	2395	4.178	−0.0
343.07	486.32	2397	4.174	−0.1

Table 2 Experimental results for the solubility of CO₂ in the ionic liquids expressed both as Henry's law constants, K_H and as mole fraction, x_2 , corrected for a partial pressure of solute of 0.1 MPa, where p is the experimental equilibrium pressure and the per cent deviation is relative to the correlations of the data reported in Table 3

T/K	$p/10^2$ Pa	$K_H/10^5$ Pa	$x_2/10^{-4}$	dev %
[N ₄₁₁₁][Ntf ₂]				
282.94	692.44	23.78	418.0	+0.2
290.17	725.19	27.93	356.1	−0.3
303.38	360.61	36.13	275.5	+0.3
303.38	779.81	36.20	274.9	+0.1
303.39	753.25	36.34	273.9	−0.3
313.24	383.51	43.33	229.8	+0.2
313.28	818.72	43.51	228.9	−0.1
323.15	405.20	51.36	194.0	+0.1
323.19	824.23	51.54	193.3	−0.1
333.09	425.91	60.28	165.3	+0.0
343.00	445.54	69.95	142.5	−0.0
343.07	890.12	70.02	142.4	−0.0
[C _{4mim}][Ntf ₂]				
283.36	675.30	23.91	415.7	+0.4
293.25	715.64	29.95	332.1	−0.2
303.21	752.47	36.41	273.3	−0.2
303.23	958.23	36.57	272.2	−0.6
303.26	933.69	36.45	273.1	−0.2
313.36	1001.18	43.51	228.9	−0.3
313.39	971.94	43.23	230.3	+0.4
323.22	820.01	50.76	196.3	−0.3
323.48	1042.44	51.08	195.0	−0.5
323.61	1008.86	50.19	198.5	+1.4
333.70	1082.50	59.09	168.6	−0.8
333.76	1044.78	57.69	172.7	+1.7
343.78	1120.07	66.52	149.9	−0.1
343.79	1080.57	67.01	148.8	−0.8
[C _{2mim}][Ntf ₂]				
283.43	650.50	24.88	399.6	+0.2
293.39	691.74	31.06	320.3	−0.2
303.39	729.85	37.90	262.6	−0.1
323.22	799.32	53.75	185.3	+0.3
343.07	863.61	72.99	136.6	−0.1

The Henry's law constants obtained from the experimental data were correlated as a function of temperature by an empirical equation of the type:

$$\ln[K_H/10^5 \text{ Pa}] = \sum_{i=0}^n A_i (T/K)^{-i}. \quad (7)$$

Table 3 Parameters of Eq. 7 used to smooth the experimental K_H results from Tables 1 and 2, along with the per cent average absolute deviation of the fit (AAD)

Gas	A_0	A_1	A_2	AAD %
[N ₄₁₁₁][Ntf ₂]				
H ₂	+18.26	-6.372×10^3	$+9.027 \times 10^5$	0.2
CO ₂	+8.769	-1.394×10^3	-5.364×10^4	0.2
[C ₄ mim][Ntf ₂]				
H ₂	+21.10	-7.674×10^3	$+1.054 \times 10^6$	0.3
CO ₂	+5.006	$+8.455 \times 10^2$	-3.863×10^5	0.6
[C ₂ mim][Ntf ₂]				
H ₂	+21.24	-7.668×10^3	$+1.046 \times 10^6$	0.2
CO ₂	+7.935	-8.376×10^2	-1.417×10^5	0.2

Fig. 1 Mole fraction gas solubilities of CO₂ (upper plot) and H₂ (lower plot) at 0.1 MPa partial pressure of the solute as a function of temperature in:
□, [N₄₁₁₁][Ntf₂];
○, [C₄mim][Ntf₂];
△, [C₂mim][Ntf₂]. Lines represent smoothed data using the parameters in Table 3

Fig. 2 Deviations from the literature values of the correlation of the Henry's law constants obtained in this work for carbon dioxide in $[\text{C}_2\text{mim}][\text{Ntf}_2]$ (*upper plot*) and in $[\text{C}_4\text{mim}][\text{Ntf}_2]$ (*lower plot*): $\square$, this work, $\blacksquare$, Cadena et al., 2004 [1] and Anthony et al., 2005 [7]; $\circ$, Kim et al., 2005 [6]; $\triangle$, Morgan et al., 2005 [8]; $\blacktriangledown$, Camper et al., 2005 and 2006 [3, 4]; $\bullet$, Lee and Outcalt, 2006 [9]; $\blacktriangle$, Baltus et al., 2004 [2]

The A_i coefficients obtained are listed in Table 3 together with the average absolute deviations obtained for each solute. These values can be regarded as an estimation of the precision of the experimental data, which in the present case is always better than 1%.

In Fig. 1 are illustrated the solubility data, expressed in mole fraction, for carbon dioxide and hydrogen in the selected ionic liquids as a function of temperature. As it can be observed, carbon dioxide is two orders of magnitude more soluble than hydrogen (mole-fraction solubilities of 10^{-2} and 10^{-4} , respectively). The solubility of the two gases in the three ionic liquids decreases with increasing temperature in the range covered by this work.

The solubility of carbon dioxide in $[\text{C}_2\text{mim}][\text{Ntf}_2]$ has been determined by different research groups, often as a function of temperature, using different experimental techniques. It is then possible to compare some of the present results with those of other authors, the comparison providing guidance to estimate the accuracy of the data in this work. A deviation plot showing differences between the various literature sources and the correlation of the present data is represented in the upper plot of Fig. 2. As it can be observed, only two sets of values, those reported by Kim et al. [6] and by Morgan et al. [8], exhibit a difference of more than 10% from the present data. The values reported by Cadena et al. [1] and by Camper et al. [3] deviate from the present work by less than 5%. In the lower plot of Fig. 2, a similar deviation plot illustrates the solubility of CO_2 in $[\text{C}_4\text{mim}][\text{Ntf}_2]$. For this ionic liquid, the

Table 4 Thermodynamic functions of solvation for H₂ in the three ionic liquids at several discrete temperatures between 283 and 343 K. $\Delta_{\text{solv}}G^\infty$ is the Gibbs energy of solvation, $\Delta_{\text{solv}}H^\infty$ the enthalpy of solvation and $\Delta_{\text{solv}}S^\infty$ the entropy of solvation. The values are consistent with $p^\circ = 0.1$ MPa

T/K	$\Delta_{\text{solv}}G^\infty/\text{kJ}\cdot\text{mol}^{-1}$	$\Delta_{\text{solv}}H^\infty/\text{kJ}\cdot\text{mol}^{-1}$	$\Delta_{\text{solv}}S^\infty/\text{J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$
[N₄₁₁₁][Ntf₂]			
283	16.51	+0.06	−58.1
293	17.13	−1.75	−64.4
303	17.80	−3.44	−70.1
313	18.53	−5.02	−75.2
323	19.30	−6.51	−79.9
333	20.12	−7.90	−84.2
343	20.99	−9.22	−88.0
[C₄mim][Ntf₂]			
283	16.82	−1.86	−66.0
293	17.52	−3.98	−73.4
303	18.29	−5.95	−80.0
313	19.12	−7.80	−86.0
323	20.00	−9.54	−91.5
333	20.94	−11.2	−96.4
343	21.93	−12.7	−101
[C₂mim][Ntf₂]			
283	16.97	−2.28	−68.0
293	17.68	−4.37	−75.3
303	18.47	−6.33	−81.9
313	19.32	−8.17	−87.8
323	20.23	−9.89	−93.2
333	21.18	−11.5	−98.2
343	22.19	−13.0	−103

values of Cadena et al. [1] agree with the ones obtained in the present work to within 5%, but the solubilities reported by both Baltus et al. [2] and Lee and Outcalt [9] deviate by more than 10% from the ones measured here. After a careful analysis of the present data, and in light of the comparisons made, we consider that the present values are precise to within $\pm 1\%$ and have an accuracy of better than $\pm 5\%$.

The variation with temperature of the solubility for the two gases studied in the three ionic liquids, expressed in terms of the Henry's law constant, is directly related to the thermodynamic properties of solvation which, in the case of gaseous solutes at low pressures, are practically identical to the thermodynamic properties of the solution [17]. The Gibbs energy of solvation is given by

$$\Delta_{\text{solv}}G^\infty = RT \ln(K_{\text{H}}/p^\circ) \quad (8)$$

where p° is the standard state pressure. The partial molar differences in enthalpy and entropy between the two states can be obtained by calculating the corresponding partial derivatives

Table 5 Thermodynamic functions of solvation for CO₂ in the three ionic liquids at several discrete temperatures between 283 and 343 K. $\Delta_{\text{solv}}G^\infty$ is the Gibbs energy of solvation, $\Delta_{\text{solv}}H^\infty$ the enthalpy of solvation and $\Delta_{\text{solv}}S^\infty$ the entropy of solvation. The values are consistent with $p^0 = 0.1$ MPa

T/K	$\Delta_{\text{solv}}G^\infty/\text{kJ}\cdot\text{mol}^{-1}$	$\Delta_{\text{solv}}H^\infty/\text{kJ}\cdot\text{mol}^{-1}$	$\Delta_{\text{solv}}S^\infty/\text{J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$
[N₄₁₁₁][Ntf₂]			
283	7.464	−14.7	−78.5
293	8.247	−14.6	−78.1
303	9.026	−14.5	−77.8
313	9.802	−14.4	−77.5
323	10.58	−14.4	−77.2
333	11.35	−14.3	−76.9
343	12.11	−14.2	−76.7
[C₄mim][Ntf₂]			
283	7.460	−15.67	−81.7
293	8.264	−14.90	−79.0
303	9.042	−14.17	−76.6
313	9.797	−13.49	−74.4
323	10.53	−12.86	−72.4
333	11.25	−12.3	−70.6
343	11.94	−11.7	−68.9
[C₂mim][Ntf₂]			
283	7.543	−15.3	−80.7
293	8.345	−15.0	−79.7
303	9.137	−14.7	−78.8
313	9.921	−14.5	−78.0
323	10.70	−14.3	−77.3
333	11.47	−14.0	−76.6
343	12.23	−13.8	−76.0

of the Gibbs energy with respect to temperature

$$\Delta_{\text{solv}}H^\infty = -T^2\partial/\partial T(\Delta_{\text{solv}}G^\infty/T) = -RT^2\partial/\partial T[\ln(K_{\text{H}}/p^0)] \quad (9)$$

$$\Delta_{\text{solv}}S^\infty = (\Delta_{\text{solv}}H^\infty - \Delta_{\text{solv}}G^\infty)/T = -RT\partial/\partial T[\ln(K_{\text{H}}/p^0)] - R\ln(K_{\text{H}}/p^0). \quad (10)$$

The values of the Gibbs energy, enthalpy and entropy of solvation are given in Tables 4 and 5 for the two gases in the three ionic liquids at seven discrete temperatures between 283 and 343 K.

The Gibbs energy of solvation is depicted in Fig. 3 for carbon dioxide (upper plot) and for hydrogen (lower plot) in the ionic liquids studied. As it can be observed, the Gibbs energy of solvation increases with increasing temperature for both gases in a similar manner for all the ionic liquids studied. This variation leads to negative enthalpies of solvation, indicating an exothermal solvation process. In the temperature range studied, and as can be observed in Fig. 4, the enthalpy of solvation is more negative for carbon dioxide than for hydrogen. For the first gas, it does not vary significantly with temperature but, in the

Fig. 3 Gibbs energy of solvation of CO₂ (*upper plot*) and H₂ (*lower plot*) at 0.1 MPa partial pressure of the solute as a function of temperature in:

- , [N₄₁₁₁][Ntf₂];
○, [C₄mim][Ntf₂];
▽, [C₂mim][Ntf₂]

case of hydrogen, it decreases significantly with temperature. For the three ionic liquids studied, the enthalpy of solvation for hydrogen approaches zero for the lower temperatures studied.

The entropy of solvation in Fig. 4, is noticeably more negative for the imidazolium based ionic liquids and varies between -60 and $-100 \text{ J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$. For all of the ionic liquids, its variation with temperature is more important for hydrogen than for carbon dioxide in the temperature range covered by this work.

Because the solubility data are sufficiently precise, the thermodynamic properties of solvation can be used to make inferences about the molecular mechanisms pertaining to the solvation of the gases in the three ionic liquids. These properties provide valuable information both about the solute–solvent interactions and about the molecular structure of the solutions: the enthalpy of solution is closely related with the cross gas–ionic liquid molecular interactions, and the entropy of solvation gives indications about the structure of the solvent molecules surrounding the solute.

Fig. 4 Enthalpy of solvation (*left*) and entropy of solvation (*right*) of the gases in ionic liquids based in the [Ntf₂] anion as a function of temperature: □, CO₂; ○, H₂

4 Conclusions

We report the solubility of carbon dioxide and of hydrogen, as a function of temperature, in three ionic liquids sharing the bis(trifluoromethylsulfonyl)amide anion. The results obtained here can be compared with values published in the literature by different research groups. These comparisons allow the estimation of a realistic value for the accuracy of the present

data. Together with a careful analysis of the sources and order of magnitude of the errors associated with our experimental technique, we consider that the accuracy of all the values presented in this paper is $\pm 5\%$.

Changing the cation of the ionic liquid has only a minor effect on the solubility of carbon dioxide. It is nevertheless noticeable that this gas is slightly less soluble in $[\text{C}_2\text{mim}][\text{NTf}_2]$. For hydrogen, its solubility is also similar in the three ionic liquids, with the differences in solubility being similar to the ones found for carbon dioxide. The relative order of the hydrogen solubility is the same as this gas is more soluble in $[\text{N}_{4111}][\text{Ntf}_2]$ followed by $[\text{C}_4\text{mim}][\text{NTf}_2]$. It can be concluded that the effect of the ionic liquid cation on the solubility of both hydrogen and carbon dioxide is similar.

The data obtained here make it possible to calculate and analyze the thermodynamic properties of solvation, which allow us to assess the molecular interactions in solution. It was observed that, although the order of magnitude of the properties of solvation is similar, their variation with temperature is quite different, pointing to probable differences in the mechanisms of solvation for the two gases studied. These differences have already been noticed for other ionic liquids [12, 13, 18].

The behavior of each gas in the presence of the imidazolium based ionic liquids is very similar. For $[\text{N}_{4111}][\text{Ntf}_2]$ as solvent, it is observed that the solvation of hydrogen is less exothermal than in the imidazolium-based ionic solvents. The entropies of solvation, of hydrogen and carbon dioxide are also less negative for $[\text{N}_{4111}][\text{Ntf}_2]$. The fact that the enthalpies of solvation are more negative for the imidazolium-based ionic liquids is explained by the existence of stronger solute-solvent interactions that are dominated by specific electrostatic forces [19].

References

1. Cadena, C., Anthony, J.L., Shah, J.K., Morrow, T.I., Brennecke, J.F., Maginn, E.J.: Why is CO_2 so soluble in imidazolium-based ionic liquids? *J. Am. Chem. Soc.* **126**, 5300–5308 (2004)
2. Baltus, R.E., Culbertson, B.H., Dai, S., Luo, H., DePaoli, D.W.: Low-pressure solubility of carbon dioxide in room-temperature ionic liquids measured with a quartz crystal microbalance. *J. Phys. Chem. B* **108**, 721–727 (2004)
3. Camper, D., Becker, C., Koval, C., Noble, R.: Low pressure hydrocarbon solubility in room temperature ionic liquids containing imidazolium rings interpreted using regular solution theory. *Ind. Eng. Chem. Res.* **44**, 1928–1933 (2005)
4. Camper, D., Becker, C., Koval, C., Noble, R.: Diffusion and solubility measurements in room temperature ionic liquids. *Ind. Eng. Chem. Res.* **45**, 445–450 (2006)
5. Dyson, P.J., Laurency, G., Ohlin, C.A., Vallance, J., Welton, T.: Determination of hydrogen concentration in ionic liquids and the effect (or lack of) on rates of hydrogenation. *Chem. Commun.* 2418–2419 (2003)
6. Kim, Y.S., Choi, W.Y., Jang, J.H., Yoo, K.-P., Lee, C.S.: Solubility measurement and prediction of carbon dioxide in ionic liquids. *Fluid Phase Equilib.* **228–229**, 439–445 (2005)
7. Anthony, J.L., Anderson, J.L., Maginn, E.J., Brennecke, J.F.: Anion effects on gas solubility in ionic liquids. *J. Phys. Chem. B* **109**, 6366–6374 (2005)
8. Morgan, D., Ferguson, L., Scovazzo, P.: Diffusivities of gases in room-temperature ionic liquids: Data and correlations obtained using a lag-time technique. *Ind. Eng. Chem. Res.* **44**, 4815–4823 (2005)
9. Lee, B.-C., Outcalt, S.L.: Solubilities of gases in the ionic liquid 1-*n*-butyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide. *J. Chem. Eng. Data* **51**, 892–897 (2006)
10. Husson-Borg, P., Majer, V., Costa Gomes, M.F.: Solubilities of oxygen and carbon dioxide in butyl methyl imidazolium tetrafluoroborate as a function of temperature and at pressures close to atmospheric pressure. *J. Chem. Eng. Data* **48**, 480–485 (2003)
11. Costa Gomes, M.F., Husson, P., Jacquemin, J., Majer, V.: Interactions of gases with ionic liquids: experimental approach. In: Rogers, R.D., Seddon, K.R. (eds.) *ACS Symp. Series Ionic Liquids III: Fundamentals, Progress, Challenges, and Opportunities*, Chap. 16, pp. 207–218. American Chemical Society, Washington (2005)

12. Jacquemin, J., Costa Gomes, M.F., Husson, P., Majer, V.: Solubility of carbon dioxide, methane, methane, oxygen, nitrogen, hydrogen, argon, and carbon monoxide in 1-butyl-3-methylimidazolium tetrafluoroborate between temperatures 283 K and 343 K and at pressures close to atmospheric. *J. Chem. Thermodyn.* **38**, 490–502 (2006)
13. Jacquemin, J., Costa Gomes, M.F., Husson, P., Majer, V.: Low-pressure solubilities and thermodynamics of solvation of eight gases in 1-butyl-3-methylimidazolium hexafluorophosphate. *Fluid Phase Equilib.* **240**, 87–95 (2006)
14. IUPAC Commission on Atomic Weights and Isotopic Abundances: Atomic weights of the elements 1985. *Pure Appl. Chem.* **58**, 1677–1692 (1986)
15. Dymond, J.H., Smith, E.B.: *The Virial Coefficients of Pure Gases and Mixtures*. Clarendon, Oxford (1980)
16. Jacquemin, J., Husson, P., Padua, A.A.H., Majer, V.: Density and viscosity of several pure and water-saturated ionic liquids. *Green Chem.* **8**, 172–180 (2006)
17. Hildebrand, J.H., Prausnitz, J.M., Scott, R.L.: *Regular and Related Solutions*. Reinhold, New York (1970), pp. 111–141
18. Costa Gomes, M.F.: Low-pressure solubility and thermodynamics of solvation of carbon dioxide, ethane, and hydrogen in 1-hexyl-3-methylimidazolium bis(trifluoromethylsulfonyl)amide between temperatures of 283 and 343 K. *J. Chem. Eng. Data* **52**, 472–475 (2007)
19. Deschamps, J., Costa Gomes, M.F., Padua, A.A.H.: Molecular simulation study of interactions of carbon dioxide and water with ionic liquids. *Chem. Phys. Chem.* **5**, 1049–1052 (2004)